

### KAPTSOV, N. A.

"Electrical Discharges in Gases and Their Use in Technology" (Elektricheskiye razryady v gazakh i ikh primeneniye v tekhnike), "Pravda," 1949, 24 pp.

A Kenteransa

**一种电影和第四次。100%型影光形流音型四型数型影响** 

WARRAN CO.		
KAPTSOV, N. A.,	Prof	PA 35/49T28
	A COLOR	
	# 1	THE FEET HOFF
	60 R 610 k	USSE, WELD BELL BY BY BELL BY BY BELL BY
		THE TO SEE TO BE
	Siecti Clig	
-	ω S LF	
	Sep 11.	P. D. S. C.
	CSR/Electricity of electrons and attend 11 Sep 48.	Electricity Corona Discil Electrons, I Lics of Elect Vacuum, "Pro oscow State trichestvo" trichestvo" sively surve ns of basic acuums. Dis ntary discha
		6 5 1 C 4 1 C 4 2 2 4 4 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
		Begin No U
	One	
	in gas	Electricity Corona Discharges Electrons, Motion sics of Electrical Di Vacuum, "Prof M. A. loscow State U imeni trichestvo" No 1 - p. lesively surveys contents ons of basic phenoment acuums. Discusses on ntary discharge procentary discharge procentary
	그 됐나 하고 하고 하는데 모든 그릇이다.	P C C C P C C P C P C P C P C P C P C P
•		
	- 호텔 : 이번 경험 : 11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	on on the one of the o
<i>4</i> − <i>4</i> −	Jan 49 Contal Co	USSR/Electricity Corona Discharges Electrons, Motion  "Thysics of Electrical Discharges in Cases and in High Vacuum," Frof N. A. Kaptsov, Dr Physicomath Sci, Moscov State U imeni Lomonosov, 16 pp  "Elektrichestvo" No 1 - p. 17-32  Extensively surveys contemporary physical representations of basic phenomena during discharge in gase elementary discharge processes on electrode surface lementary processes in gaseous space, and movement \$5/49126
		# 4 4 6 7 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
	- 1000년 전 1000년 시간 시간 시간 시간 시간 시간 (1000년 100년 100년 100년 100년 100년 100년 10	E of the Apr
		90000
		Gases Physi 16 pp 1cal 1cal charg 3h ga strod,
	그렇게 하기를 보고 한 닭이들로 하겠다니다.	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
	35))	
	Sub-	Jan 49 and in someth cometh in gas es surfac movemen
	7.5	4 4 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
:		USSR/Electricity  Corona Discharges  Electrons, Motion  "Physics of Electrical Discharges in Gases and in High Vacuum," Prof M. A. Kaptsov, Dr Physicomath Sci, Moscov State U imeni Lomenosov, 16 pp  "Elektrichestvo" No 1 - p. 17-32  Extensively surveys contemporary physical representations of basic phenomena during discharge in gases, elementary discharge processes on electrode surfaces elementary processes in gaseous space, and movement  35/49728
	는 발생 이 동생 등록 보면 사용한다고 불리하는 및 및 호텔스트리와, 그는 원리한 발음 모든 생활 하는 발생 :	
от в селинествення в нешти переменення в нешто при нешто не при нешто в досто в 1900 године. Под 1900 године в При при нешто в нешти переменення в нешто при нешто при нешто при нешто под 1900 године.	an and the substitution of the section of the substitution of the	nations: an intercupant and off the section of the contract of

KAPTSOV, N. A.

"Petr Nikolayevich Lebedev (1866 - 1912")", 39 pp, 1950.

KAPTSOV, N. A.

Tibis: Stertric Pasnomana in Cases and Vacuum

Author: Kaptsov, A. A.

Mother Organization:

Israing Agercy: State Publishing House of Technical and Theoretical

数据数据转换的 60万元的 2000年的 100元代码 120元代码 120元元代码 100元元代码 100元元代码 100元元代码 100元元代码 100元元代码 100元元代码 100元元代码 100元元代码 100元

Literature

Iccation: Foscow-Leningred

Pates of Issue: 1950 , 836 pp.

Holdings--DLC::

ATIC 26604-1

Holdings-Other Libraries

Contents: Theromionic and auto-electronic (cold) emissions

External photo-effect

fonization and excitation of gas particles in non-exprasible collissions of the first and second types.

Radiation of a gaseous discharge

Townseni discharge and the conversion of a discharge from

dependent to independent character.

CIA-RDP86-00513R000720510020-4" APPROVED FOR RELEASE: 06/13/2000

KAPTSOV, N. A. (editor); LEB, L.

"Basic Process of Electric Discharges in Gases" (Csnovnyye protsessy elektricheskikh razryadov v gazakh), Gosudarstvennoye Izdatel'stvo Tekhnikoteoreticheskoy Literatury, 672 pp, 1950.

Book W-22459, 22 Apr 52

KAPTSOV N. A.

181739

USSR/Electricity - Gas Lamps, High-Pressure Apr 51

"High-Pressure Lamps," N. A. Kaptsov, D. A. Goukhberg

"Uspekhi Fiz Nauk" Vol XLIII, No 4, pp 620-662

Reviews high-pressure mercury vapor illuminating lamps and describes constr, characteristics and application of high-pressure inert gas-filled lamps.

181739

- 1. KAPTSOV, N. A.
- 2. USSR (600)
- 4. Electronics
- 7. "Electronics." Reviewed by N. A. Kaptsov. Sov.kniga. No. 9, 1952.

9. Monthly List of Russian Accessions, Library of Congress, January 1953, Unclassified.

KAPTSOV, N.A.

Physicists

Recollections of Petr Nikolayevich Lebedev. Usp. fiz. nauk 46 no. 3, 1952

Monthly List of Russian Accessions, Library of Congress, August 1952. Unclassified.

SAMPLE STREET STREET STREET STREET STREET STREET STREET

KAPTSOV, N. A.

"Electronics", Gostekhizdat, Moscow, 1953 - 468 pb.

The name of the book is somewhat misleading as it does not appear to be a general textbook on the subject of electronics, but deals with, more particularly, the theory of electronic discharge phenomena in vacuum and gases. The book was intended for the specific purpose as a text for advanced students in radio physics.

Translation summary - XXII - 3

KAPTSOV, N.A.; PETROV, Vasiliy Vladimirovich.

150th anniversary of the appearance of V.V.Petrov's book "News of galvano-voltaic experiments." Usp.fiz.nauk 50 no.2:303-307 Je '53. (MLRA 6:7) (Petrov, Vasilii Vladimirovich, 1761-1834) (Electricity)

APPROVED FOR RELEASE: 06/13/2000 CIA-RDP86-00513R000720510020-4"

[Electronics] Elektronika. M, Gostekhizdat, 1954. (MLRA 8:5)
(Electronics)

KAPTSOV, N.A.

FD-742

USSR/Physics - Electric gas discharge

Card 1/1

: Pub 146-12/22

Author

: Kaptsov, N. A., and Popov, N. A.

Title

: Flash of electric discharge in gases on alternating current of audio

frequency in tubes with external and internal electrodes.

Periodical

: Zhur. eksp. i teor. fiz., 27, 97-102, Jul 1954

Abstract

: Flash and discharge glow in tubes with external and internal electrodes is analyzed in relation to frequencies of applied voltage. Using external electrodes an unstable discharge was found at low voltages and a stable one at high voltages. At a frequency of 10 kc curves of flash voltage vs. frequency tend to overlap. 7 foreign

references.

Institution

: Moscow State University

Submitted

November 28, 1953

CIA-RDP86-00513R000720510020-4" **APPROVED FOR RELEASE: 06/13/2000** 

KAptSOV N.A.

USSR/Physics - Vacuum technology

Card 1/1 : Pub. 86 - 4/40

Authors : Kaptsov, N. A., Prof.

Title : The technology of high vacuums

Periodical : Priroda 43/4, 33-44, Apr 1954

Abstract: The term high vacuum is explained as a rarefication of sufficiently high degree as to permit a molecule to traverse the length of a container without colliding with another molecule. The electrical uses of vacuums are enumerated. A description is given of an oil-filled rotary pump which will produce a vacuum of less than 0.001 mm of mercury atmospheric pressure. For higher vacuums this pump serves for the first stage of rarefication and a mercury pump, a description of which is also given, reduces the atmospheric pressure further. Methods for preserving a vacuum and measuring the degree of rarefication are explained. Illustrations; diagrams; drawings; graph.

Institution:

Submitted: .... Translation M-3.053,44,

KAPTSOV, N. A. Prof.

-10 +4 May 15

"Nature of High-Frequency Discharge," a paper delivered at the Section of Radiophysics, Physics Faculty, Moscow University, Conference on Radiophysics, Moscow State University, Vest. Mosk. Universitet, Ser. Fiz-Mat. i Yest. Nauk, No.6, 1955

Sum. 900, 26 Apr 56

USSR/Physics - Spectrum of crypton Card 1/1 50 V, N. A. Pub. 129-4/20

FD-2164

Author

: Devyatov, A. M., and Kaptsov, N. A.

Title

A STATE OF THE STA Investigation of the excitation functions of certain spectral lines of

krypton

Periodical: Vest. Mosk. un., Ser. fizikomat. i yest. nauk, 10, No 2, 27-36, Mar 1955

Abstract

: Up to the present time the excitation functions of the energy levels and spectral lines of a small number of elements have been investigated; namely H, He, Ne, Ar, Hg, Zn, Cd, Na, etc. (1927-1952). In the present work the authors determine the relative functions of excitation of certain spectral lines of krypton by an optical method. They describe the procedure of the experiment and experimental arrangement; the results obtained are shown in 17 graphs (excitation function for various Kr lines and wave lengths). Fifteen references; e.g. four by B. M. Yavorskiy (1944-1947); A. N. Zaydel', V. K, Prokof'yev, and S. N. Rayskiy, Tablitsy spektral'nykh liniy (Tables of spectral lines), GITL, Moscow-Leningrad, 1952.

Institution:

Submitted

: September 4, 1954

KAPTSOV, N.A., professor, (Moskva)

Ivan Filippovich Usagin. Fiz. v shkole 15 no.5:90-91 S-0 '55. (Usagin, Ivan Filippovich, 1855-) (MIRA 9:1)

KAPTSOV, H.A.; professor, doktor fisiko-matematicheskikh nauk.

Electric discharges in gases. Nauka i shizn' 22 no.5:9-11 My '55 (MIRA 8:6)

1. Moskovskiy gosudarstvennyy universitet imeni M.V. Lomonosova. (Electric discharges through gases)

SIDOROV, Mikhail Alekseyevich; KAPTSOV, H.A., professor, redaktor; MEZENTSEV, V.A., redaktor; AKHLAMAOV, S.N., tekhnicheskiy redaktor

[From shavings to electricity] Ot luchiny do elektrichestva. Pod red. N.A.Kaptsova. IEd. 2-oe. Noskva, Gos. iEd-vo tekhniko-teoret. lit-ry, 1956. 61 p. (Nauchno-populiarnaia biblioteka, no.56)
(Lighting) (MIRA 9:9)

KAPISON, NIKOLAY A.

Call Nr: AF 1119832

AUTHOR:

Kaptsov, Nikolay A., Moscow State University

TITLE:

Electronics (Elektronika)

PUB. DATA:

State Publishing House of Technical and Theoretical Literature, Moscow, 1956, 2d ed., 459 pp., 20,000 copies

ORIG. AGENCY:

None given.

EDITORS:

Alekseyev, D.M. and Murashova, N.Ya., Reviewer:

Spivak, G.V., Prof.

PURPOSE:

Approved by the Ministry of Higher Education of the USSR as a textbook for students of State Universities. The present 2nd edition of the book is based on the

author's earlier lectures at the Radio-Physics


Department of Moscow University.


COVERAGE:

See Table of Contents

<del>(a. 1-1/1-)</del>







VASIL'YEVA, M.Ya.; KAPTSOV, M.A.

Studying the difference between the igniting and extinguishing voltage of glow discharge under various conditions. Vest. Nosk. un. 11 no.2:29-35 F '56. (MLRA 9:8)

1. Kafedra elektroniki.

(Electric discharges)

# POPOV, N.A.; KAPTSOV N.A.

Investigation of an intermediate frequency discharge occurring between high frequency and low audio frequency discharges at low gas pressure. Zhur.eksp.i teer. fiz. 30 no.1:68-76 Ja 156. (MIRA 9:7)

1. Heskevskiy gesudarstvennyy universitet. (Electric discharges through gases)

是**的第三人称形式的现在分词**的复数形式的一种,这种一种的一种,这种种种的一种,是一种种的一种,是一种的一种的一种,是一种的一种的一种,是一种的一种的一种的一种,

KAPTSOV, Nikolay Aleksandrovich, prof.; PLONSKIY, A.F., red.; MURASHOVA, N.Ya., tekhn.red.

[Pavel Nikolaevich IAblochkov; his life and work] Pavel Nikolaevich IAblochkov; ego zhizn' i deiatel'nost', Moskva, Gos.izd-vo tekhniko-teoret. lit-ry, 1957. 95 p. (MIRA 10:12) (IAblochkov, Pavel Nikolaevich, 1847-1894)

KAPTSOV, N. A.

"High Frequency and Ultra High Frequency Discharges in Gases."

PRINT

paper presented at Second All-Union Conference on Gaseous Electronics, Moscow, 2-6 October '58.

24(3) AUTHORS: Kuzovnikov, A.A., Kaptsov, N.A.

**医眼睛表现的形式的形式的影响的影响的影响,这些影响的大型,但是是这种形成的影响,那些影响,这些地域的影响的影响,而是不知识的影响的影响的影响。**"你吃了,"

SOV/155-58-5-27/37

TITLE:

Discharge Power and the Character of the Discharge Current

for Frequencies of 1.5 up to 9 mc

PERIODICAL:

Nauchnyye doklady vysshey shkoly. Fiziko-matematicheskiye

nauki,1958,Nr 5,pp 158-166 (USSR)

ABSTRACT:

With the aid of the experimental equipment described in [Ref 1 7 the discharge power as well as the magnitude and character of the discharge current were measured in the given frequency interval. The discharge arising in the air between a sphere and a plane under atmospheric or lower pressure was investigated. An approximative theory of the appearance is proposed. Among others it is stated : The power necessary for maintaining the discharge increases with increasing frequency of the external electric field. The transition from the corona discharge to the torch takes place under equality of the amplitudes of the active and reactive components of the electron current. An approximative investigation of the directed electron motion is possible, if it is based on the solution of the equation of motion of the averaged electron

Card 1/2

27

Discharge Power and the Character of the SOV/155-58-5-27/37 Discharge Current for Frequencies of 1.5 up to 9 mc

in the electric field under consideration of the coefficient of friction and of the frequency of the natural oscillations of the electrons. The properties of these solutions show good qualitative coincidence with experimental results. In the corona discharge the directed electron motion is stronger than the disordered motion caused by heat; in the torch it is inverse. There are 5 figures, 1 table, and 16 references, 10 of which are Soviet, 4 American, and 2 German.
Tsyan'Gao Yun, Candidate, and N.N. Bulatova are mentioned.

ASSOCIATION: Moskovskiy gosudarstvennyy universitet imeni M.V.Lomonosova (Moscow State University imeni M.V. Lomonosov)

SUBMITTED: June 16, 1958

Card 2/2

THE RESERVE

9(0) AUTHOR:

Kaptsov, N. A., Professor

507/30-59-8-2/56

TITLE:

Gas Electronics - a Topical Field of Physics

PERIODICAL:

Vestnik Akademii nauk SSSR, 1959, Nr 8, pp 12 - 17 (USSR)

ABSTRACT:

The theoretical bases of gas electronics have not yet been worked out. No clear explanations are available concerning the formation of electric discharges in gases. There is no quantitative theory of the formation and propagation of streamers. The problems of the development of high-tension discharges are of special importance. The most powerful ones are produced and investigated under laboratory conditions at voltages of several million volt between the electrodes. The phenomena of ball lightning, electric arc and the discharge at high and superhigh frequency are still to be investigated. The investigation of the state of ionized gas, called gas plasma, is considered to be the most essential problem of gas electronics from the scientific point of view, existing in modern technics. A distinguen is made between isothermal and non-isothermal plasma. The formation of electromagnetic oscillations is characteristic of plasma; in this connection

Card 1/2

Gas Electronics - a Topical Field of Physics

sov/30-59-8-2/56

a distinction is made between electron and ion oscillations. Plasma has also magnetic properties. Gas in form of plasma is at present used for the solution of important technical problems. For the purpose of investigating the plasma properties it is necessary to find its parameters under various conditions. However, the solution of this task is possible only on the basis of the joint experimental and theoretical work of a number of physical and technical laboratories and institutes.

Card 2/2

KAPTSOV, N.A.

P.N.Lehedev and his school. Trudy Inst.ist.est.i tekh. 28: 106-110 159. (MIRA 13:5) (Lebedev, Petr Nikolaevich, 1866-1912)

### PHASE I BOOK EXPLOITATION

sov/4705

Padiofizicheskaya elektronika (Radiophysical Electronics)[Moscow]Izd-vo Mosk. univ., 1960. 561 p. Errata slip inserted. 15,000 copies printed.

Ed.: N. A. Kaptsov, Professor; Tech. Ed.: M. S. Yermakov.

PURPOSE: This book has been approved by the Ministry of Higher and Secondary Special Education, USSR, as a textbook for schools of higher education. It can be also used by scientific personnel working in the fields of radio engineering and electronics.

COVERAGE: The book presents problems of vacuum, cathode, semiconductor, and gas electronics, on which is based the operation of vacuum-tube and gas-filled devices, including microwave devices and also apparatus and instruments used in electron optics. It is assumed that the readers of this book havea preliminary preparation in the fundamentals of nuclear physics, quantum mechanics, statistical physics and electrodynamics. The book was written by a group of lecturers of the Physics Division of Moscow State University.

Card 1/10

# APPROVED FOR RELEASE CO6/13/2000 CIA-RDP86-00513R660 #20510020-4"

Chapters I, II, and III were written by Professor N. A. Kaptsov; Ch. IV. by Professor S. D. Gvozdover and Docent V. M. Lopukhin; Ch. V. by Professor G. V. Spivak and Assistant Ye. M. Dubinina; Ch. VII. by Docent A. A. Zaytsev and Professor N. A. Kaptsov; Ch. VIII. by Professor N. A. Kaptsov and Assistant G. S. Solntsev. The authors thank Professor S. Yu. Luk'yanov and Docent M.D. Karasev , who reviewed the book. There are 76 references: 68 Soviet (including 14 translations), 6 English, and 2 German.

### TABLE OF CONTENTS:

#### Foreword

1. 2. 3.	Subject of Physical Electronics. High-Vacuum Electronics Introduction Transmission of electric current through a high vacuum Space charges in gaseous, liquid and solid media Physics of electron tubes	9 13 21 21
5• 6.	Semiconductor Electronics Electron energy levels in crystals Impurity semiconductors	29 29 32
	Law of electron distribution along the separate energy levels in semiconductor energy bands Density of energy states in any energy band of a crystal	36 39
Card-2		

Radiop	physical Electronics	sov/4705
9•	Computation of level position of electrochemical potenti semiconductors	als in
10.	Dependence of the electric conductivity of semiconductor temperature	s on 48
11.	Photoconductivity of crystals and its use in photoresist	
12.		54
13.	Photoeffect of the barrier layer	56
14.	Semiconductor (dry) rectifiers	56 63
	Modern types of dry rectifiers	71
16.	Crystal detectors	
17.	Transistors	73 76
Ch. II	I. Cathode Electronics	85
18.	Subject of cathode electronics	
-	Laws of thermionic emission	85 86
20.	Effect of the external field on a cathode surface during thermionic emission	
	MEINTONIC CHIRDION	95
Card.3	/10	

Radiophysical Electronics	80V/4705
21. Effect on the work function of thin layers of foreign so on the cathode surface. Film cathodes	ubs <b>tance</b> s 98
22. Surface ionization	100
23. Thermionic semiconductor emission	102
24. Thermion distribution according to energy levels	104
25. Oxide-coated cathodes	104
26. Spot field	113
27. Various designs of hot cathodes	115
28. Cold (autoelectronic) emission	115
29. Positive ion emission from the plate (thermo-ionic emiss	•
30. Extrinsic photoeffect	125
31. Methods of experimental investigation of metal and semi- conductor photoeffects	<b>-</b> 129
32. Normal and selective photoeffects. Effect of thin surface on the photoeffect of metals	
33. Fowler's theory. Dependence of the photoeffect on metal	
temperature	132
34. Theory of extrinsic photoeffect	137
35. Special features of photoelectron emission from semicond	
cathodes	139
Card # 10	

Radiophy	vsical Electronics SOV/4	1705
36.	Cesium-oxide photocathodes	142
37.	Antimony-cesium cathodes	145
38.		
7	occurring in complex photocathodes	147
39•	and the contract of the contra	148
40.	Abnormal secondary emission. Malter effect	156
41.		158
		159
43.		163
1. 1.	and excited and neutral atoms	164
<del>////</del> *		165
45.	Shot effect and flicker effect	10)
en. Tit	Missesses Pleatwenies	168
	Microwave Electronics	168
40.	Introduction	200
47.	Differential equation systems of electrodynamics and	174
	microwave electronics	-17
48.	Excitation of resonators by electron flows. Concept of	180
	induced current	182
49•	Double-cavity klystron	105
Card 5/	<del>- 10</del> .	

Radioph	ysical Electronics Sov/4705	
80. 81. 82. 83.	Statistical discharge lag and time of discharge formation Streamer theory of gas breakdown Starting the discharge in long tubes High-vacuum breakdown	386 388 399 400 401
85.	Preconduction current pulses  Devices based on the use of dependent discharges, preconduction pulses, and starting of independent discharges. Tonization chambers and counters  Arresters and trigatrons	
87. 88. 89. 90. 91.	Cas-discharge plasma Methods of determining plasma parameters Plasma oscillations Distribution of plasma electrons according to energy levels Theory of a homogeneous positive column Striated positive column Deionization of plasma Ionospheric layers and their role in radio wave propagation	409 409 411 425 431 440 449 453 456
Card 87	<del>(20</del>	

Radiophy	ysical Electronics SOV/4705	
Ch. VIII	I. Various Types of Electric Discharges in Gases and Their Role in Radio and Other Technical Fields	461
	Boundary condition of electric discharges in gases	461
96.	Glow discharge	462
97•	Normal and abnormal fall in cathode potential during a glow discharge	466
98.	Cathode sputtering	468
99.	Glow-discharge devices	470
100.	Various types of arc discharge. Arc discharge with a thermionic cathode (artificially heated)	472
101.	Electric arc	474
102.	Theory of electric-arc ribbon	479
103.	Behavior of electric-arc ribbon in super-high power current	
	pulses	483
104.	Mercury-arc rectifiers and ignitrons	487
105.	Gas-filled tube rectifiers and thyratrons. Arc rectifiers	490
	Use of an electric arc as a source of high temperature	494
107.	Electric arc and other types of arc discharge as a source of light	t495
	Spark discharge and lightning	499
Card-9/	10	

109.	Corona discharge. Corona-discharge field		507
110.	Bipolar corona discharge. Brush discharge. Initial corona pulses		515
111.	Application of corona discharge in engineering		516
	H-f discharges and their role in radio engineering.		517
113.	Special features of directed and random electron motion	in an	
	h-f discharge field		518
114.	Various types of h-f discharges		534
115.	H-f pulse discharge		542
116.	Diffusion theory of h-f discharge ignition		544
117.	Applications of h-f discharges		550
Bibliogr	raphy		555
Subject	Index		558
AVATLABI	E: Library of Congress	JP/rsm/fal 1-25-61	

Card 10/10

88015

26.2311 24,2120 (1049, 1160,1482) S/139/60/000/006/009/032 E073/E335

AUTHORS:

Kuzovnikov, A.A. and Kaptsov, N.A.

TITLE:

Investigation of a High-frequency Discharge in the

Range Between 1.5 and 15 Mc/s. III

Izvestiya vysshikh uchebnykh zavedeniy, PERIODICAL: Fizika, 1960, No. 6, pp. 64 - 70

TEXT: The mechanism of development of a high-frequency corona discharge and its change to a torch discharge cannot be studied solely on the basis of the theory of unbounded uniform plasma (Ref. 1). On the basis of experimental data, published earlier by the authors (Refs. 2, 6, 7), they suggest a mechanism of the development of such a discharge which is based on the conceptions of the avalanche-streamer theory. Application of the ideas of the avalanche-streamer theory to the high-frequency corona discharge at atmospheric and subatmospheric (300 - 400 mm Hg) pressures can be justified by the fact that both in the steady-state and in surge corona as well as in high-frequency corona individual localised discharge

Card 1/4

880k5 S/139/60/000/006/009/032 E073/E314

Investigation of a High-frequency Discharge in the Range Between 1.5 and 15 Mc/s. III

canals can be observed. The characteristics of the highfrequency corona (Ref. 2) are analogous to those of the steady-state (Refs. 3, 4) and surge (Ref. 5) corona discharges. In the earlier work of the authors (Refs. 2, 6, 7) it is shown that on increasing the voltage the high-frequency corona passes successively through the following three main stages (Ref. 2): 1) in the initial stage the discharge is in the form of fine channels which are distributed fanlike on the corona producing electrode; 2) in the second stage a bright central canal and numerous clearly visible side canals form which penetrate deep into the discharge gap; 3) in the third stage a high-frequency arc forms. mechanism of development of a high-frequency corona discharge was investigated for the frequencies 1.5, 2, 3.7, 4, 6.5 and 8.7 Mc/s. The authors conclude that the mechanism of development of high-frequency corona discharges can be

Card 2/4

88045 \$/139/60/000/006/009/032

Investigation of a High-frequency Discharge in the Range Between 1.5 and 15 Mc/s. III

elucidated on the basis of the avalanche-streamer theory. In the initial stage of the corona and the torch discharge individual, short-length, rectilinear discharge canals form as a result of successive superposition on each other of electron avalanches and also as a result of development of an avalanche canal during oscillatory movement of the electrons under the effect of the high-frequency field. If the voltage amplitude increases to a certain value the formation of streamers in the corona discharge becomes possible. The discharge canals, which can be seen with the naked eye during this stage of the corona, are formed as a result of reforming of the streamer canal, as a result of secondary processes on the temporary cathode or as a result of oscillatory movement of the electrons under the effect of the high-frequency field. Under the given conditions streamer formations and consequently also the formation of individual visible canals of the high-frequency corona occur in the discharge at

E073/E335

Card 3/4.

88025 S/139/60/000/006/009/032 E073/E335

Investigation of a High-frequency Discharge in the Range Between 1.5 and 15 Mc/s. III

atmospheric pressure if the active duration of the half-cycle of the voltage is equal to or greater than 0.03 uses. The torch discharge is a high-frequency plasma which is formed during numerous half-cycles of the high-frequency field and is drawn out upwards by the convection currents of the air. There are 1 table and 16 references: 11 Soviet and 5 non-Soviet.

ASSOCIATION:

Moskovskiy gosuniversitet imeni M.V. Lomonosova

(Moscow State University imeni M.V. Lomonosov)

SUBMITTED:

October 21, 1959

Card 4/4

KAPTSOV, N.A.

Petr Nikolaevich Lebedev's role in the training of young scientists. Usp.fiz.nauk 77 no.4:582-588 Ag '62.

(MIRA 15:8)

(Physics—Study and teaching)
(Lebedev, Petr Nikolaevich, 1866-1912)

LEBEDEV, Petr Nikolayevich, akademik; KRAVTS, T.P., red. (1866-1912);
KAPTSOV. N.A., prof., red.; YELISEYEV, A.A., dots., red.;
Endauaul, r.u., red. izd-va; Makuni, Ye.V., tekhn. red.


[Sollected works] Sobranie sochinenii. Moskva, Izd-vo AN SSSR, 1963. 434 p. (MIRA 16:9)

1. Chlen-korrespondent AN SSSR (for Kravts).

(Lebedev, Petr Nikolaevich, 1866-1912) (Physics)

VVEDENSKIY, B.A., glav. red.; VUL, B.M., glav. red.; SHTEYNMAN, R.Ya., zam. glav. red.; BALDIN, A.M., red.; VONGOVSKIY, S.V., red.; GALANIN, M.D., red.; ZERLOV, D.V., red.; ISHLINSKIY, A.Yu., red.; KAPITSA, P.L., red.; KAPTSOV, N.A., red.; KOZODAYEV, M.S., red.; LEVICH, V.G., red.; LOYTSYANSKIY, L.G., red.; LUK'YANOV, S.Yu., red.; MALYSHEV, V.I., red.; MIGULIN, V.V., red.; REBINDER, P.A., red.; SYRKIN, Ya.K., red.; TARG, S.M., red.; TYABLIKOV, S.V., red.; FEYNBERG, Ye.L., red.; KHAYKIN, S.E., red.; SHUBNIKOV, A.V., red.

[Encyclopedic physics dictionary] Fizicheskii entsiklopedicheskii slovar'. Moskva, Sovetskaia Entsiklopediia. Vol.4. 1965. 592 p. (MIRA 18:1)



#### "APPROVED FOR RELEASE: 06/13/2000 CIA-RDP86-00513R000720510020-4 AND THE PERSON OF THE PROPERTY OF THE PERSON OF THE PERSON

KAPTSOV

USSR/Chemical Technology - Chemical Products and Their Application. Industrial Organic Synthesis

I-1

Abs Jour

: Ref Zhur - Khimiya, No 1, 1958, 2154

Author

: Topchiyev, A.V., Kaptsov, N.N.

Inst

: Academy of Sciences USSR

Title

: Utilization of Nitrogen Oxide for the Nitration of Paraf-

Orig Pub

: Sb.: Khim. pererabotka neft. uglevodorodov. M., AN SSSR,

Abstract

: A study was made of the effect of temperature and space velocity on the course of the reaction of concurrent interaction of NO, O2 and n-pentane. In the investigation use was made of a reactor with a reaction zone enclosed at the same time by a cold and a hot wall. Temperature of the preheater was varied in the range of 430-540°, that of the

Card 1/2

USSR/Chemical Technology - Chemical Products and Their Application. Industrial Organic Synthesis.

I-l

Abs Jour

: Ref Zhur - Khimiya, No 1, 1958, 2154

rection zone within 290-338°, and space velocity was varied from 0.93 to 2.30 min<sup>-1</sup>. The molar ratios n-C<sub>2</sub>H<sub>12</sub>: NO: 0, were from 1: 0.49: 0.44 to 1: 0.97: 0.6. Extent of conversion of  $C_2$ H<sub>12</sub> and NO and yield of nitroparaffins reach maximum values at a temperature of the preheater of 445° and a space velocity somewhat above 1.0, and are of 17, 20 and 22¢, respectively. The authors assume that the reaction between NO, 02 and  $C_2$ H<sub>12</sub> takes place within a certain zpne that is intermediate between the hot and the cold wall of the reactor. A diagram of a laboratory unit for the nitration of paraffins is included.

Card 2/2

KAPTSOV,

USSR/Organic Chemistry. Theoretical and General

E-1

Questions of Organic Chemistry.

Abs Jour

Ref Zhur - Khimiya, No. 8, 1957, 26659.

Author

Inst

Topchiyev, A.V.; Kaptsov, N.N. Academy of Sciences of USSR. Inst. Petroleum Primary Radical Formation in Reaction

Title

of Vapor Phase Nitrating of Alkanes by

Nitrogen Dioxide.

Orig Pub

Izv. AN SSSR, Otd. khim. n., 1956, No. 7,

863 - 868.

Abstract

One of the surmised reactions at the vapor phase nitrating of alkanes by NO2 is the formation of alkyl radicals according to the equation  $RH + NO_2 \rightarrow R + HNO_3$  (1). The computations show that the change of the free energy  $\triangle z^0$  at 25° is 26.6 kcal/mol at the reaction of CH4 with NO2, Azgos is

Card 1/2

S/020/60/132/02/35/067 B011/B002

AUTHORS:

والمسيد بيناي

Topchiyev, A. V., Academician, Kaptsov, N. N., Zalesskaya, L. N.

TITLE:

Nitration of Parsoxydiphenyldimethylmethane Acetate in the Presence

of Urea

PERIODICAL: Doklady Akademii nauk SSSR, 1960, Vol. 132, No. 2, pp. 371-373

TEXT: The authors proved that during the nitration of paraoxydiphenyldimethylmethane acetate, one of the three nitro groups enters a non-phenolic cycle of the molecule (see scheme). For the purpose of purification p-oxydiphenyldimethylmethane (ODDM) (commercial by-product of the phenol acetone production) was first recrystallized from a mixture of benzene-petroleum ether. The ODDM crystals are white, needle-shaped and have their melting point at 73°-75°. Production of the acetate: ODDM was dissolved in an aqueous KOH solution with an addition of ethanol, and 180 g of acetic anhydride were quickly added. After it was cooled down for half an hour by adding lumps of ice, or when the mixture was put on ice, the solution separated in layers. It was extracted by means of ether. When the ether was distilled off, the remaining substance was a colorless, ether. When the ether was distilled almost without decomposition at 327° at

Card 1/3

Nitration of Paraoxydiphenyldimethylmethane Acetate in the Presence of Urea

s/020/60/132/02/35/067 B011/B002

atmospheric pressure. The melting point of this acetate was  $180^{\circ}-182^{\circ}/1.5$  mm. The molecular weight was determined to be 250 and calculated to be 254. The acetate easily dissolves in benzene, benzine, o-xylene, and other solvents. Nitration of the acetate by means of HNO3 at 150-200 leads to the formation of picric acid. This can be prevented if the acetate is poured off at lower temperatures and if the reaction mass is left standing at a lower temperature. Thus low yields of a yellowish crystalline substance develop with a melting point of 127°. It was analyzed to be the trinitro derivative of p-ODDM. Its molecular weight was determined to be 356 and calculated to be 347. In order to avoid the oxidative action of HNO3, the authors nitrated ODDM acetate in the presence of urea. Table 1 shows that in this case, the trinitro compound develops with a considerably higher yield. Urea however, must be added after the acetate has been poured off, otherwise only picric acid would develop. The position of the nitro groups was proven by oxidation with chromic acid. A small amount (0.07 g) of a solid yellow substance was obtained with a melting point between 238° and 241°. The authors compared it with para-nitro-benzoic acid whose melting point is at 241° (Scheme). There are 1 table and 4 references, 1 of which is Soviet.

ENERGY DE PROPERTO DE LA PROPERTO D

Card 2/3

11

35526 s/020/62/143/003/024/029 B101/B144

15.0050

11.2215

Topchiyev, A. V., Academician, Kaptsov, N. N., Kalyuzhnaya,

G. D., Mityayeva, A. I., and Balitskaya, I. Ye.

TITLE:

AUTHORS:

Interaction of polymers and copolymers of 2-methyl-5-vinyl

pyridine with aromatic nitro compounds

PERIODICAL: Akademiya nauk SSSR. Doklady, v. 143, no. 3, 1962, 621 - 624

TEXT: To test the activity of the pyridine-nitrogen atom in addition reactions, polymers (PI) of 2-methyl-5-vinyl pyridine (I) and its styrene copolymers (SI) were reacted with various polar compounds. A PI with softening point 186°C and three SI with I: styrene ratio of 5:1,3:1, and 1 : 1 were used. To test the effect of basicity on the reaction with dinitro compounds, the SI with ratio 1: 1 was nitrated by means of 73% HNO3 and 24% H<sub>2</sub>SO<sub>4</sub> at 20°C (decomposition of this nitro compound occurred above 200°C). 2.5%, 5%, and 10% solutions were prepared from PI and SI in a mixture 1: 1 of dinitro toluene (DNT) and dinitro xylene (DNX); their viscosity was measured and was found to increase with length of The same behavior was found in the case of nitrated SI. An heating. Card ·1/3

S/020/62/143/003/024/029 B101/B144

Interaction of polymers...

extraction of PI dissolved in DNT + DNX by means of benzene was unsuccessful. The increasingly dark red and finally dark brown polymer became impoluble insoluble in benzene, and its melting point was higher than 250°C. From this, cross linking was concluded, and the structure

was proposed. As unpurified DNT + DNX mixture caused a considerable

Card 2/4

5/204/63/003/001/008/013 E075/E436

Topchiyev, A.V. (deceased), Kusakov, M.M., AUTHORS:

Kalyuzhnaya, G.D., Kaptsov, N.N., Koshevnik, A.Yu.,

Razumovskaya, E.A.

Characterization of the properties of homo- and TITLE:

copolymers of 2-methyl-5-vinylpyridine by the methods

of light scattering and viscosimetry

PERIODICAL: Neftekhimiya, v.3, no.1, 1963, 90-93

The authors determined the molecular weights and other properties of polymerized 2-methyl-5-vinylpyridine and its TEXT: 1:1 copolymer with styrene. The polymerizations were carried out by heating 2-methyl-5-vinylpyridine at 80°C for 12 hours in glass ampules with 0.1% benzoylperoxide. From the light scattering and viscosimetry data the following relationship was obtained

$$[\eta] = 6.17 \times 10^{-4} M_W^{0.615}$$

where  $[\eta]$  - intrinsic viscosity and  $M_W$  - mean molecular weight. The mean molecular weights of the polymer fractions obtained by Card 1/2

S/204/63/003/001/008/013 E075/E436

Characterization of ...

petroleum-ether precipitation, ranged from  $1 \times 10^6$  to  $3 \times 10^4$ . The mean molecular weights of the copolymer were  $4.3 \times 10^5$  and  $1.1 \times 10^5$  for the polymerization times of 12 and 6 hours respectively. There is 1 table.

ASSOCIATION: Institut neftekhimicheskogo sinteza AN SSSR (Institute of Petrochemical Synthesis AS USSR)

SUBMITTED: August 18, 1962

Card 2/2

KAPTSOV, N.P., dots.; KRYLOV, A.V., dots., otv. red.

[Complex movement of a point; methodological textbook on theoretical mechanics] Slozhnoe dvizhenie tochki; uchebnometodicheskoe posobie po teoreticheskoi mekhanike. Itv. red. A.V. Krylov, Moskva, Mosk. in-t neftekhim. i gazovoi promyshl. im. I.M.Gubkina, 1959. 17 p. (MIRA 15:2)

AUTHOR: Kaptsova, I.N. TITLE:-The Use of Gravimeters in Underground Mining Galleries, for SOV/132-58-11-10/17 the Study of Mineral Depostis (O primenenii gravimetrov v podzemnykh gornykh vyrabotkakh dlya izucheniya rudnykh me-PERIODICAL: Razvedka i okhrana nedr, 1958, Nr 11, pp 36 - 40 (USSR) ABSTRACT: The Kafedra gravimetrii i nebesnoy mekhaniki Gosudarstvennogo Astronomicheskogo instituta imeni P.K. Shternberga (the Chair of Gravimetry and of Celestial Mechanics of the State Astronomical Institute imeni P.K. Shternberg ) organized an experimental gravimeter survey of copper or deposit, both from the surface and from an underground gallery. The gravimeter GAK-3M was used. The experiment showed that the joint interpretation of both operations made the solution of the reversed problem of the gravimeter survey more accurate. The experiment and the solution of the problem are given in detail. There are 3 graphs, 1 table and 7 references, 4 of which are ASSOCIATION: (MGU. GAISh)

Card 1/1


APPROVED FOR RELEASE: 06/13/2000 CIA-RDP86-00513R000720510020-4"

MARENNIKOVA, S.S.; KAPTSOVA, T.I.

Age-dependence of susceptibility of white mice to variola virus. Acta virol. (Praha) [Eng] 9 no.3:230-234 My'65.

1. The Moscow Scientific Research Institute of Viral Preparations, Moscow, U.S.S.R.

APPROVED FOR RELEASE: 06/13/2000 CIA-RDP86-00513R000720510020-4"



AUTHORS:

Delimarskiy, Yu. K., Kaptsova, T. N.

SOV/78-3-12-23/36

TITLE:

Polarographic Investigation of a Solution of Titanium Dioxide in Molten Sodium Metaphosphate (Polyarograficheskoye issledovaniye rastvora dvuokisi titana v rasplavlennom metafosfate natriya)

PERIODICAL:

Zhurnal neorganicheskoy khimii, 1958, Vol 3, Nr 12,

pp 2751-2756 (USSR)

ABSTRACT:

In the present paper a solution of titanium dioxide in molten sodium metaphosphate was investigated polarographically using solid stationary electrodes. The linear dependence between N and  $\mathbf{i}_{d}$  was expressed by means of the following equation:

 $i_d = kN$  (1). In the polarogram two waves appear, which indicate the step-wise reduction of the titanium (IV) ion. The reduction apparently occurs in the following steps:

Ti<sup>4+</sup> + e → Ti<sup>3+</sup>,
Ti<sup>3+</sup> + 3e → Ti.

The polarographic waves plotted for the system under investigation correspond to the equation of Geyrovskiy-Il¹kovich. The half-wave potential  $\rm E_{1/2}$  is independent of the concentration.

Card 1/2

SOV/78-3-12-23/36

Polarographic Investigation of a Solution of Titanium Dioxide in Molten Sodium Metaphosphate


A linear dependence exists between E and lg  $\frac{i}{i_d-i}$  . The energy

of activation of the diffusion current for the first and second wave were determined. The activation energy of the first wave varies from 8.6 to 19.2 kcal/mol and the second from 20.5 to 21.4 kcal/mol. There are 5 figures, 3 tables, and 19 references, 10 of which are Soviet.

SUBMITTED:

September 30, 1957

Card 2/2



ACCESSION NR: AP3003994  AUTHORS: Kaptsova, T. N.; Delimarskiy, Yu. K.  TITLE: Folarographic analysis of vanadium, molybdenum, tungsten, and iron oxides fused with scdium methaphosphate   SCURCE: Ukrainskiy khimicheskiy zhurnal, v. 29, no. 7, 1963, 714-722  TOPIC TAGS: germanium, vanadium, tungsten, molybdenum, polarography, iron, sodium ABSTRACT: This study is a continuation of a previous polarographic study of metal oxides in a fused media of sodium metaphosphate. The present study is made of germanium takes place in a stepwise procedure with subsequent formation of mides. At low concentrations of GeO2, only one break is observed. It was established that the oxides of vanadium, tungsten and molybdenum are reduced only lished by polarographic analysis. The reduction of iron oxide takes place in two ographic maximums are explained by the depolarizing action of the adsorbed procurrent was calculated for all studied oxides. Orig. art. has: 3 tables,	ACCESSION NR:	EWP(q)/EWT(m)/BDS	AFFTC/ASD	JD/WH	
TITLE: Polarographic analysis of varadium, molybdenum, tungsten, and iron oxides fused with sodium methaphosphate \( \text{N} \) \( \text{N}	.15 1			/0073/63/029/	007/0714/0722
TITLE: Polarographic analysis of varadium, molybdenum, tungsten, and iron oxides fused with sodium methaphosphate \( \text{N} \) \( \text{N}	AUTHORS: Kapts	OVA. T. N.: Delimenski.	V. w		الماعان المعاددات
fused with scdium methaphosphate \( \text{National variation}, \text{ molybdenum, tungsten, and iron oxides} \)  SCURCE: Ukrainskiy khimicheskiy zhurnal, v. 29, no. 7, 1963, 714-722  TOPIC TAGS: germanium, vanadium, tungsten, molybdenum, polarography, iron, sodium ABSTRACT: This study is a continuation of a previous polarographic study of metal oxides in a fused media of sodium metaphosphate. The present study is made of GeO2, V2O5, MoO3, WO3, and Fe2O3 oxides in the same media. The reduction of germanium takes place in a stepwise procedure with subsequent formation of phosphides. At low concentrations of GeO2, only one break is observed. It was established that the oxides of vanadium, tungsten and molybdenum are reduced only lished by polarographic analysis. The reduction of iron oxide takes place in two obraphic maximums are explained by the depolarizing action of the adean decits of electrolysis are explained by the depolarizing action of the odean decits of electrolysis are explained by the depolarizing action of the odean decits of electrolysis are explained by the depolarizing action of the odean decits of electrolysis are explained by the depolarizing action of the selectrolysis are explained by the depolarizing action of the selectrolysis are explained by the depolarizing action of the selectrolysis are explained by the depolarizing action of the selectrolysis are explained by the depolarizing action of the selectrolysis are explained to the selectrolysis action of the selectrolysis are explained to the selectrolysis action of the selectrolysis are explained to the selectrolysis action of th			240 44		<b>デ</b> カー
SCURCE: Ukrainskiy khimicheskiy zhurnal, v. 29, no. 7, 1963, 714-722  TOPIC TAGS: germanium, vanadium, tungsten, molybdenum, polarography, iron, sodium ABSTRACT: This study is a continuation of a previous polarographic study of metal oxides in a fused media of sodium metaphosphate. The present study is made of germanium takes place in a stepwise procedure with subsequent formation of phosphides. At low concentrations of GeO2, only one break is observed. It was established that the oxides of vanadium, tungsten and molybdenum are reduced only lished by polarographic analysis. The reduction of iron oxide takes place in two obraphic maximums are explained by the depolarizing action of the obtained polar-ducts of electrolysis are explained by the depolarizing action of the oderated	TITLE: Polaro	graphic analysis of vans	idium, molyba		60
TOPIC TAGS: germanium, vanadium, tungsten, molybdenum, polarography, iron, sodium ABSTRACT: This study is a continuation of a previous polarographic study of metal oxides in a fused media of sodium metaphosphate. The present study is made of germanium takes place in a stepwise procedure with subsequent formation of phides. At low concentrations of GeO <sub>2</sub> , only one break is observed. It was established that the oxides of vanadium, tungsten and molybdenum are reduced only lished by polarographic analysis. The reduction of iron oxide takes place in two ographic maximums are explained by the depolarizing action of the advantal dicts of electrolysis and molybdenum are reduced only	Tused with sodi	methaphosphate 0	201,000	tungste	n, and iron oxides
ABSTRACT: This study is a continuation of a previous polarographic study of metal oxides in a fused media of sodium metaphosphate. The present study is made of germanium takes place in a stepwise procedure with subsequent formation of phosphides. At low concentrations of GeO <sub>2</sub> , only one break is observed. It was established that the oxides of vanadium, tungsten and molybdenum are reduced only lished by polarographic analysis. The reduction of iron oxide takes place in two observed. Their half-wave potentials are close to each other. The obtained polar-ducts of electrolysis are the description of the address description of the address description of the address description of the address description of electrolysis are explained by the depolarizing action of the address description.	SCURCE: Themas	<b>(1)</b>	V (	27 1	1 nn
ABSTRACT: This study is a continuation of a previous polarographic study of metal oxides in a fused media of sodium metaphosphate. The present study is made of germanium takes place in a stepwise procedure with subsequent formation of phosphides. At low concentrations of GeO <sub>2</sub> , only one break is observed. It was established that the oxides of vanadium, tungsten and molybdenum are reduced only lished by polarographic analysis. The reduction of iron oxide takes place in two observed. Their half-wave potentials are close to each other. The obtained polar-ducts of electrolysis are the description of the address description of the address description of the address description of the address description of electrolysis are explained by the depolarizing action of the address description.	ovisi	iskly knimicheskiy zhurn	al, v. 29, no	0. 7. 1963. 7	14.700
oxides in a fused media of sodium metaphosphate. The present study of metal GeO2, V2O5, MoO3, WO3, and Fe2O3 oxides in the same media. The reduction of germanium takes place in a stepwise procedure with subsequent formation of phides. At low concentrations of GeO2, only one break is observed. It was established that the oxides of vanadium, tungsten and molybdenum are reduced only lished by polarographic analysis. The reduction of iron oxide takes place in two ownship maximums are explained by the depolarizing action of the obtained polar-ducts of electrolysis are explained by the depolarizing action of the oderated	TOPIC TAGS: CO.			1, 4,00,	14-155
oxides in a fused media of sodium metaphosphate. The present study of metal GeO2, V2O5, MoO3, WO3, and Fe2O3 oxides in the same media. The reduction of germanium takes place in a stepwise procedure with subsequent formation of phides. At low concentrations of GeO2, only one break is observed. It was established that the oxides of vanadium, tungsten and molybdenum are reduced only lished by polarographic analysis. The reduction of iron oxide takes place in two ownship maximums are explained by the depolarizing action of the obtained polar-ducts of electrolysis are explained by the depolarizing action of the oderated	8	marrium, vanadium, tungs	ten, molybder	um, polarogra	inhy inon assert
germanium takes place in a stepwise procedure with subsequent formation of phides. At low concentrations of GeO <sub>2</sub> , only one break is observed. It was established that the oxides of vanadium, tungsten and molybdenum are reduced only lished by polarographic analysis. The reduction of iron oxide takes place in two ographic maximums are explained by the depolarizing action of the obtained polar-ddicts of electrolysis are explained by the depolarizing action of the adventure.	473577771077				Land & Tront & Souling
germanium takes place in a stepwise procedure with subsequent formation of phides. At low concentrations of GeO <sub>2</sub> , only one break is observed. It was established that the oxides of vanadium, tungsten and molybdenum are reduced only lished by polarographic analysis. The reduction of iron oxide takes place in two ographic maximums are explained by the depolarizing action of the obtained polar-ddicts of electrolysis are explained by the depolarizing action of the adventure.		ETHINA SALE LAND			
phides. At low concentrations of GeO <sub>2</sub> , only one break is observed. It was established that the oxides of vanadium, tungsten and molybdenum are reduced only their trivalent state during the electrolysis in the phosphate bath as established by polarographic analysis. The reduction of iron oxide takes place in two ographic maximums are explained by the depolarizing action of the obtained polar-ducts of electrolysis.	oxides in a fuse	study is a continuation	of a previou	s polarograni	lic etudu ac
tablished that the oxides of vanadium, tungsten and molybdenum are reduced only their trivalent state during the electrolysis in the phosphate bath as established by polarographic analysis. The reduction of iron oxide takes place in two ographic maximums are explained by the depolarizing action of the obtained polar-ducts of electrolysis.	oxides in a fuse GeO2, V2O2, MOO2	study is a continuation d media of sodium metapl	of a previou hosphate. Th	s polarograph	nic study of metal
to their trivalent state during the electrolysis in the phosphate bath as established by polarographic analysis. The reduction of iron oxide takes place in two ographic maximums are explained by the depolarizing action of the adaptation of the ad	oxides in a fuse GeO <sub>2</sub> , V <sub>2</sub> O <sub>5</sub> , MoO <sub>3</sub> germanium takes	study is a continuation d media of sodium metapl , WO3, and Fe2O3 oxides	of a previou hosphate. The in the same	s polarograph present stumedia. The	nic study of metal
lished by polarographic analysis. The reduction of iron oxide takes place in two ographic maximums are explained by the depolarizing action of the advantage polar-	oxides in a fuse GeO <sub>2</sub> , V <sub>2</sub> O <sub>5</sub> , MoO <sub>3</sub> germanium takes phides. At low	study is a continuation d media of sodium metapl , WOz, and Fe <sub>2</sub> O <sub>3</sub> oxides place in a stepwise proc	of a previous hosphate. The in the same cedure with s	s polarograph e present stu media. The r ubsequent for	nic study of metal dy is made of reduction of
stages. Their half-wave potentials are close to each other. The obtained polar- dects of electrolysis are explained by the depolarizing action of the advent-	oxides in a fuse GeO <sub>2</sub> , V <sub>2</sub> O <sub>5</sub> , MoO <sub>3</sub> germanium takes mhides. At low tablished that t	study is a continuation d media of sodium metaple, WO <sub>3</sub> , and Fe <sub>2</sub> O <sub>3</sub> exides place in a stepwise proceeding of GeO <sub>2</sub> ,	of a previous hosphate. The in the same cedure with a only one breath	s polarographe present students. The rubsequent for sky is observed.	nic study of metal dy is made of eduction of mation of phos-
of raphic maximums are explained by the depolarizing action of the adverted	oxides in a fuse GeO <sub>2</sub> , V <sub>2</sub> O <sub>5</sub> , MoO <sub>3</sub> germanium takes phides. At low tablished that to their trivale	study is a continuation d media of sodium metaple, WO <sub>3</sub> , and Fe <sub>2</sub> O <sub>3</sub> exides place in a stepwise proceeding of GeO <sub>2</sub> , the exides of vanadium, the state of vanadium of va	of a previous hosphate. The in the same cedure with some only one breatungsten and in the control of the contro	s polarograph e present atomedia. The rubsequent for ak is observe	nic study of metal dy is made of eduction of mation of phos-
dicts of electrolysis at the depolarizing action of the adapt and	oxides in a fuse GeO2, V2O5, MoO3 germanium takes phides. At low tablished that to their trivale lished by polaro	study is a continuation of media of sodium metaple, WO3, and Fe <sub>2</sub> O <sub>3</sub> oxides place in a stepwise processor of GeO <sub>2</sub> , he oxides of vanadium, that state during the electronsists.	of a previous hosphate. The in the same cedure with some only one breatungsten and attrolysis in	s polarograph e present stumedia. The rubsequent for ak is observe molybdenum ar	nic study of metal dy is made of eduction of mation of phos-d. It was es-
	oxides in a fuse GeO <sub>2</sub> , V <sub>2</sub> O <sub>5</sub> , MoO <sub>3</sub> germanium takes phides. At low tablished that to their trivale lished by polaro stages. Their he	study is a continuation of media of sodium metaple, WO3, and Fe2O3 oxides place in a stepwise processor of GeO2, he oxides of vanadium, that state during the electronic analysis. The results	of a previous hosphate. The in the same cedure with a conly one breatingsten and a ctrolysis in reduction of	s polarographe present attended as The rubsequent for ak is observe molybdenum ar the phosphete iron oride to	nic study of metal dy is made of eduction of mation of phos-d. It was es-e reduced only bath as estab-
ard 1/2 calculated for all studied oxides. Orig. art. had 7	oxides in a fuse GeO <sub>2</sub> , V <sub>2</sub> O <sub>5</sub> , MoO <sub>2</sub> germanium takes phides. At low tablished that to their trivale: lished by polarousinges. Their he of the content o	study is a continuation of media of sodium metaple, WO3, and Fe2O3 oxides place in a stepwise processor concentrations of GeO2, he oxides of vanadium, that state during the electronic analysis. The raif-wave potentials are	of a previous hosphate. The in the same cedure with a conly one breatungsten and attrolysis in reduction of close to each	s polarographe present students. The rubsequent for ak is observe molybdenum ar the phosphate iron oxide tan other. The	nic study of metal dy is made of eduction of mation of phos-d. It was es-e reduced only bath as estab-kes place in two
	oxides in a fuse GeO <sub>2</sub> , V <sub>2</sub> O <sub>5</sub> , MoO <sub>2</sub> germanium takes phides. At low tablished that to their trivale: lished by polaro, syages. Their he ographic maximum ducts of electron	study is a continuation of media of sodium metaple, WO3, and Fe2O3 oxides place in a stepwise processor of GeO2, he oxides of vanadium, that state during the electronic oxides. The raif-wave potentials are state explained by the decree of the state of	of a previous hosphate. The in the same cedure with a conly one breatingsten and corrolysis in reduction of close to each epolarizing a	s polarograph of present students. The rubsequent for ak is observe molybdenum arthe phosphate iron oxide tan other. The action of the	nic study of metal dy is made of reduction of mation of phos-d. It was es-e reduced only bath as estab-kes place in two obtained polar-

	L 17703-65 ACCESSION	NR: AP300399		The second secon	To To		ر در
		and 9 formules	그 성격하는 제공하는 하나요?				1
	ASSOCTATIO	No Traddaul s					T = 0
	general an	d inorganic che	eschey i neorgan	dcheskoy khir	mii AN Vkrs	SR (Instit	ute of
8 . 1 Year of a 1	SUPMITTED:				Ukrssr)		
[基数 至]			DATE ACQ:	15Aug63		ENCL:	00
	SUB CODE:	CH, EL	no ref sov	: 009		CTHER: (	02
							<i>N</i> 2
装出							
****							

DELIMARSKIY, Yu.K.; ANDREYEVA, V.N.; KAPTOOVA, T.N.

Reaction of metal oxides with fused modium metaphosphate, 12v.
AN SSSR. Neorg. mat. 1 no.1:150-155 Ja 165. (MIRA 18:5)

1. Kiyevskiy tekhnologicheskiy institut pishahevey promyahlennosti.

CHKANIKOV, D.I., kand.sel'skokhozyaystvennykh pauk, KAPTSYNEL', Yu. M.

Herbicidal action of aliphatic chlorinated carboxylic acids. Izv.
TSKhA no.6:80-92 '60. (MIRA 13:12)
(Acids, Fatty) (Herbicides)

是那些的是的是否的是这种研究的对象的对称的企业。这个对象的对象。12.11亿元的系统,全部的第三人称为,这个对象的现在分别的数据的对象的对象的对象的对象的对象的

# KAPTURENKO, A.M.

Introducing the economic accountability in planning organizations. Transp. stroi. 9 no.11:40-41 N '59 (MIRA 13:3)


1. Starshiy ekonomist Glavtransproyekta.
(Building research) (Construction industry--Accounting)

 REDYMENTION ARREPT-CHAINED BOOK EXPLORATION 507/3226  REDYMENSORY ARREPT-CHAINED SPICETURE DEPOSITIONS VI.  *Spreasmanys destinanty applications proistods vi.  *Configurations of the Intercollegists Scientife and Fedulationary Configuration and Fedulations and Fedulatio	۰ ۸	\ d	1 /	члс	٠,	<u>.</u>	Y <u></u>	· S	22	8	8	7	-	· <u></u>			
	I BOOK EXPLOITATION		file and ng Indus	Sponsoring Agendies: Lemingradakiy politakhnicheskiy institut im- M.I. Ealinina, Mauchno-tekhnicheskoye obshchastvo mashinotroit- ely, Latingradakoye odelaniye, and Mauchno-tekhnicheskoye obshch- estvo metallungoy, Laningradskoye odelemiye.	Resp. Ed.: V.S. Smirnov, Doctor of Technical Salences, Professor; Ed.: W.B. Paylov.	FURFOR: These proceedings of the conference are intended for apecialists in the rolling industry.	COTEMAE: The articles of this collection cover various theoretical and practical problems of rolling, such as: pressure, spread, efficiency of rolls, determination of deformation; forces required, pass design, optimus conditions for rolling, experiences of various plants, modernization of equipment, signimum-clad steel,	and rolling of bonferrous metals. We personalities are mentioned. References appear after each article. Smirnov V.5. [changradsky politekheitehesky institut im. Smirnov V.5. [changradsky politekheitehesky institut im. Kalinin]] ***T. K. Liftun [changradsky politekheitehesky institut im. K. I. Kalinin]] *********************************	Shayum, V.L. (SiOC im. Ordhonikidze, Kramatorak) 01d Kramatorak machine-Balliding Plant in the Drive for Technical Progress	. [Dasproptrov- tallurgical saure in			Starthenko, D.I. [Zhdanovskiy metallurgicheskiy institut (Zhdanov Netallurgical Institute) Forward Slip, Retardation and Spread in Molling With Normal and Extra High Drafts	ret g		a. Ealinin) ip	Klisenio, V.M. [Institut chemoy metallurgii AN UUSR (Institute of Perrous Metallurgy, AS UG. 287), Attavi) design of Reses with Staching Effect [top end bottom of pass have small tapers] and the Experimental Decemblation of Side Pressure of Work in es

Par 60 Brands Grands Ser 42 Brands Ser 42 Br	Commence of the control of the contr	CYCLONG TOLERAN AND A STATE OF THE STATE OF	skly institut	, i	1. of Publishing House: R.A. Delina; Tech.	WRPOLE: This collection of articles is intended for tecnnical and scientific personnel in matallurgy and in methanical engineering. It will also be of interest to designers of rolling equipment.	COVERMEN: This collection of articles treats the theory of rolling. It discusses such factors as the total and the unit presenter of the work on rolls, ancests of rolling, former slip, spread, etc. It also includes results obtained from investigation of real quality, rolling of quaf torn sheets and other problems.		Rolls The investigation was carried out to develop a reliable method of measuring unit pressure on the contact surface, and to obtain, by measurement, date on distribution of unit pressure during rolling with verious drafts of strips having various funful professions and widths.	Chekmarev, A.P., and P.E. Kilmenko. Experimental Investigation of Distribution of Unit Pressures on the Contact Surface During: Rolling in drooved Rolls	Chekmarev, A.P., and Rudoy, V.S. [Candidate of Technical Sciences, Institut obernoy uscallurgil AT Unions, and successionation and more leadingil AT Unions, and secondary and the secondary statement of Percous Metallurgy of the Action of Sciences of the Unrainian SSR, and the Allurgy of Sciences of the Unrainian SSR, and the Allurgy of Science for Plainial, The Contact Surface, and Fresaure an Rolla in Filger [Rockeite] Molling SSR. The authors present new actions for measuring present on Palling Discussing present of Percolling pages with 219, 273 and 225 mm disserters, and for determining the instant area of contact.	Varidin, Va.L. [Candidate of Twohnical Sciences]. Pressure on Tolis in Rotary Rolling of Tubes on a Short Mandrel The author compares experimental data on the total and unit pressures with the results obtained through wang formulas the	Delibor derived. A.P., V.M. Kilsenko, W.I., Meleshko, M.M. Saflyen, V.D. Ghebrarey, A.F., V.M. Kilsenko, W.I., Mersaure on V.D. Ghebraroy, and S.M. Rabinovich [Definer]. Fressure on Rolls in Situating Mill the section with the authority describe the methods, instruments, and results of an investigation carried out at the "Importants at with all on horizonts and vertical rolls at all rolling.	Saf'yan, M.M. [Candidate of Technical Sciences]. Experimental Investigation on the Lewer-Ara of Monents in Cold Malling.  The author describes investigation on the above subject, and gives the total pressure on Folls in Cold Tolling of seed at the total pressure on Folls in Cold Tolling of seed.	Chekamarev, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1,	<u>Butivey, M.M.</u> (Candidate of Technical Sciences). Derivation of a 152 pormula for Spread of Rolling on Elain Rolls of apread it rolling. The author presents a metrod of calculation of apread it rolling. It is based on theoretical determination of streams in the contact area in the contact area in the contact area.
		: :		<del></del>	<del>, , , ,</del> ,	·	· · · · · · · · · · · · · · · · · · ·	erre A		<del>,</del>		<del> </del>		<del></del>	<u> </u>	

"APPROVED FOR RELEASE: 06/13/2000

CIA-RDP86-00513R000720510020-4



8/137/61/000/006/026/092 A005/A101

AUTHORS:

Chekmarev, A.P., Kapturov, L.Ye., Klimenko, P.L.

TITLE:

Experimental investigation of the distribution of specific pressure

over the contact surface during rolling on smooth rolls

PERIODICAL:

Referativnyy zhurnal. Metallurgiya, no. 6, 1961, 1 abstract 6D4 ("Nauchn. tr. Dnepropetr. metallurg. in-t", 1950, no. 39, 5 - 29)

The authors substantiate a method selected for investigating speci-TEXT fic pressures with the aid of a dynamometer functioning with a glued-on wire resistance pickup of a surface subjected to stretching. The investigations were made on a laboratory two-high mill with rolls of 260 mm diameter and 350 mm length. The experimental methods are described in detail. Pb-strips of 22, 16, 10, 6, 4 and 2 mm thickness, 50, 35 and 20 mm width, and 350 mm length each, were rolled, and it was established that: 1) specific pressures are non-uniformly distributed across the deformation seat; over its length they are highest in the center and least at the edges; 2) the absolute magnitude of specific deformation decreases with a reduced width of the strip; 3) during rolling of thick strips with a reduction of \le 23% tensile stresses arise which entail a decrease

Card 1/2

Experimental investigation ...

\$/137/61/000/006/026/092 A006/A101

of specific pressure on the contact aurface. To investigate the distribution of specific pressure during non-uniform deformation, special concave and convex Presections were rolled. It was found that compressive stresses increased the specific pressure in strip sections subjected to stronger compression and that tensile stresses reduced the specific pressure in less compressed sections of the strip.

V. Pospekhov

[Abstracter's note: Complete translation]

Card 2/2

CHEKMAREV, A.P., akademik; KAPTUROV, L.Ye., inzh.

Experimental investigation of specific pressures in hot rolling.
Nauch. trudy IMI no.39:278-292 160. (MIRA 13:10)

1. AN USSR (for AN USSR).
(Rolling mills)

CHEKMAREV, A. P., akademik; KAPTUROV, L. Ye., inzh.; RABINOVICH, S. N., inzh.

Metal pressure on rolls and cogging conditions on a threehigh sheet rolling mill in the Novo-Kramatorsk machinery plant. Nauch, trudy DMI no.48:239-249 162. (MIRA 15:10)

1. Akademiya nauk Ukrainskoy SSR (for Chekmarev).

(Kramatorsk-Machinery industry)
(Rolling(Metalwork)

CHEKMAREV, A. P., akademik; RABINOVICH, S. N., inzh.; KAPTUROV,
L. Ye., inzh.

Investigating the grooving and the wear of rolls on a two-high thin sheet rolling mill. Nauch. trudy DMI no.48:250-256 162. (MIRA 15:10)

1. Akademiya nauk Ukrainskoy SSR (for Chekmarev).

(Rolls(Iron mills)). (Mechanical wear)

CHEKMAREV, A. P., akademik; RABINGVICH, S. N., inzh.; <u>KAPTUROV</u>, <u>L. Ye.</u>, inzh.; <u>MASHKIN</u>, L. F., inzh.


Automatic shape adjustment of sheet mill rolls by means of a mechanical grinding device. Nauch. trudy IMI no.48:265-274 162. (MIRA 15:10)

(Rolls(Iron mills)) (Grinding and polishing)
(Electronic control)

KAPTUROV, L. Ye., inzh.

Experimental investigation of the effect of rolling speed on specific pressure. Nauch. trudy DMI no.48:311-315 '62. (MIRA 15:10)

(Rolling(Metalwork))



KAPTUROVA, O.L.

Subject

: USSR/Chemistry

AID P - 3489

Card 1/1

Pub. 152 - 4/21

Authors

Ganz, S. N. and S. I. Kapturova

Title

A STATE OF THE PARTY OF THE PAR Kinetics of formation of nitric acid in mechanical absorbers with a large number of revolutions

Periodical

: Zhur. prikl. khim., 28, 6, 585-596, 1955

Abstract

In mechanical absorbers with a large number of revolutions the gas is thoroughly mixed with the liquid, and the oxidation of NO to NO<sub>2</sub> proceeds at a higher rate. Two tables, 13 diagrams, 12 references, all

Russian (1900-1953).

Institution : None

Submitted

: N 4, 1953

DELIMARSKIY, Yu. K.; KAPTSOVA, T. N.; BOYKO, K. M.

Polarographic investigation with fused sodium metaphosphate as the support. Ukr. khim. zhur. 28 no.5:595-599 '62. (MIRA 15:10)

1. Institut obshchey i neorganicheskoy khimii AN UkrSSR.

(Polarography) (Sodium metaphosphate)

DELIMARSKIY, Yu.K.; KAPTSOVA, T.N.

Polargraphic investigation with fused sodium metaphosphate as the support. Part 2: Polarography of copper, silver, cadmium, and lead oxides. Ukr. khim. zhur. 28 no.7:802-802 162. (MIRA 15:12)

l. Institut obshchey i neorganicheskoy khimii AN UkrSSR. (Metallic oxides) (Polarography)


CHERMAREV, A. P., akademik; KAPTUROV, L. Ye., inzh.; RABINOVICH, S. N., inzh.


Metal pressure on rolls and cogging conditions on a two-high thin sheet rolling mill. Nauch. trudy DMI no.48:257-264 162. (MIRA 15:10)


1. Akademiya nauk Ukrainskoy SSR (for Chekmarev).

(Rolling(Metalwork))

1







. KHELYHE,

137-58-1-1765

Translation from: Referativnyy zhurnal, Metallurgiya, 1958, Nr 1, p 240 (USSR)

AUTHOR: Kaptyug, L.S.

TITLE: On Woody Fracture of Steel (O shifernom izlome stali)

PERIODICAL: V sb.: Metallovedeniye, Leningrad, Sudpromgiz, 1957, pp 253-263

ABSTRACT:

A critical analysis is presented of the results of some researches on schistose woody fracture in steel, and a discussion is presented of the connection between the manifestation of schistosity and the state of the metal at the moment of fracture testing. It is shown that schistosity and exfeliation do not exist in schistose metal before fracture testing as defects characteristic of disruption of the integrity of the metal. They develop in the testing process. and only in that portion of the volume of the specimen which undergoes serious plastic deformation prior to the moment of fracture of the specimen. In steel predisposed to formation of schistosity or exfoliation, no heat treatment is capable of eliminating these defects from fibrous fracture. When fibrous or woody fracture is replaced by crystalline fracture, the appearance of schistosity is impaired or prevented, but this does not improve

Card 1/2

137-58-1-1765

On Woody Fracture of Steel

DECEMBER OF THE PROPERTY OF TH

the quality of the steel. The formation of woody fracture of steel is intimately related to contamination thereof by non-metallic inclusions, while the formation of crystalline fracture does not depend upon their presence in the steel. It is recommended that an evaluation of the quality of steel for tendency to schistosity and exfoliation be made on transverse sections, and that the testing of notched specimens to fracture be done slowly under a press.

N.K.

1. Steel -- Fracture -- Analysis

Card 2/2

KAPTYUG, I.S., kand, tekhn.nauk; SYSHCHIKOV, V.I., inzh.

Some results of testing titanium and its alloys for friction and wear. Sudostroenie 24 no.8:46-48 Ag '58. (MIRA 11:10)

(Titanium—Testing)

AUTHORS: Kaptyng, I.S. (Candidate of Technical Sciences) and Syshchikov, V.I. (Engineer)

Influence of Alloying on the Friction Properties of Titanium (Vliyaniye legirovaniya na friktsionnyye

PERIODICAL: Metallovedeniye i Termicheskaya Obrabotka Metallov, 1959, Nr 4, pp 22-27 (USSR)

ABSTRACT: The authors investigated the friction properties of some titanium alloys produced in an induction furnace from a de-gassed sponge metal containing 0.01% C, 0.14% Si, 0.16% Fe, 0.08% Mg. The smelting and the alloying were effected in graphite crucibles in an argon atmosphere. The chemical compositions and the mechanical properties of the heats are entered in Table 1 (p 23), and it can be seen that 0.38 - 0.80% C passed from the crucibles into the alloy. Ingots of 70 - 90 mm diameter were forged into rods of 15 mm diameter which were then cooled in air. From the latter, specimens were produced for tensile tests and also for friction tests. In the experiments the coefficient of friction and the tendency

Card 1/3 to seizing were investigated (at specific pressures of 10, 100 and 300 kg/cm2; at each of these 30 sliding

。 1. 1915年 - 1915年 -

Influence of Alloying on the Friction Properties of Titanium motions were made), as well as the wear resistance, the hardness and the microstructure. The obtained results are entered in tables and plotted in graphs. The authors arrived at the following conclusions: 1) Alloying of titanium brought about only a slight reduction in the static friction coefficient (from 0.55 to 0.45 in a rubbing pair with titanium and from 0.20 to 0.15 in a rubbing pair with brass) and in the depth of penetration of the damage in the case of dry sliding friction. 2) Titanium and the investigated titanium alloys proved to have a very low wear resistance against sliding friction; the wear was 15 - 30 times as high as that of brass, bronze or stainless steel. 3) The investigated titanium alloys as well as pure titanium are unsuitable for components subjected to friction under high pressure. Card 2/3 However, they can be used in rubbing pairs with brass or

Influence of Alloying on the Friction Properties of Titanium bronze in the case of relatively low loads.

There are 5 figures and 4 tables.

Card 3/3

# KAPTYURINA, Anna Dmitriyevna

[Lumbosacral radiculitis] Polasnichno-kresttsovyi radikulit.
Moskva, Medgiz, 1960. 15 p. (MIRA 13:8)

(NERVES, SPINAL--DISEASES)

KAPTYUSHIN, I., prepodavatel

Instruction maps in laboratory work. Prof.-tekh. obr. 17 no. 11:13 N '60. (MIRA 13:12)

1. Uchilishche mekhanizatsii sel'skogo khozyaystva No 32. Saratovskaya oblast'.

(Farm mechanization--Study and teaching)

KAPTYUSHIN, I., prepodavatel'

Final lesson. Prof.-tekh.obr. 19 no.1:10 Ja '62. (MIRA 15:1)

1. Borskoye uchilishche mekhanizatsii sel'skogo khozyaystva No.1, Kuybyshevskaya oblast'. (Farm mechanization--Study and teaching)

DARKANBAYEV, T.B.; KAPTYUSHINA. G.A.

Sugar and starch content of grain and flour of Kazakhstan. Izv. AN Kazakh.SSR.Ser.biol.no.10:87-93 '55. (MIRA 9:4)

1. Institut botaniki AN Karssr. (KAZATHSTAN-WHRAT)

Kazakhstan wheat grains contained reducing sugars 0.17-0.22, sucrose 2.43-3.3, and starch 50.5-66.0%; 72% yield flour contained reducing sugars 0.15-0.2, sucrose 1.76-2.25, and starch 70.41-75.63%. Generally the grain of hard wheat contained more sucrose than that of soft wheat.

KAPTYUSHINA, G.A., Cand Bio Sci-(dias/ "Biochemical indicators and breadbaking properties of certain new varieties of Kazakhstan wheat." Alma-Ata, 1958. 21 pp (Min of Higher Education USSR. Kazakh State U im S.M. Kirov), 150 copies (KL, 30-58, 125)

-48 -